arXiv:1610.05684v1 [math.NT] 15 Oct 2016

Recursive constructions of k-normal polynomials over finite fields

Mahmood Alizadeh
Department of Mathematics, Ahvaz Branch, Islamic Azad University, Ahvaz, Iran
alizadeh@iauahvaz.ac.ir

Saeid Mehrabi
Department of Mathematics, Farhangian University, Tehran, Iran
saeid_mehrabi@yahoo.com

Abstract

The paper is devoted to produce infinite sequences of k-normal polynomials $F_u(x) \in \mathbb{F}_q[x]$ of degrees np^u $(u \geq 0)$, for a suitably chosen initial k-normal polynomial $F_0(x) \in \mathbb{F}_q[x]$ of degree n over $\mathbb{F}_q$ by iteratively applying the transformation $x \to \frac{x^p - x}{x^p - x + \delta}$, where $\delta \in \mathbb{F}_q$ and $0 \leq k \leq n-1$.

Keywords:Finite fields; normal basis; k-normal element; k-normal polynomial.
Mathematics Subject Classification:12A20

1 Introduction

Let $\mathbb{F}_q$ be the Galois field of order $q = p^m$, where p is a prime and m is a natural number, and $\mathbb{F}_q^*$ be its multiplicative group. A *normal basis* of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$ is a basis of the form $N = \{\alpha, \alpha^q, ..., \alpha^{q^{n-1}}\}$, i.e. a basis that consists of the algebraic conjugates of a fixed element $\alpha \in \mathbb{F}_{q^n}^*$. Such an element $\alpha \in \mathbb{F}_{q^n}$ is said to generate a normal basis for $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$, and for convenience, is called a *normal element*.

A monic irreducible polynomial $F(x) \in \mathbb{F}_q[x]$ is called *normal polynomial* or *N-polynomial* if its roots are linearly independent over $\mathbb{F}_q$. Since the elements in a normal basis are exactly the roots of some N-polynomials, there is a canonical one-to-one correspondence between N-polynomials and normal elements. Normal bases have many applications, including coding theory, cryptography and computer algebra systems. For further details, see [9].

Recently, the k-normal elements over finite fields are defined and characterized by Huczynska et al [8]. For each $0 \leq k \leq n-1$, the element $\alpha \in \mathbb{F}_{q^n}$ is called a k-normal element if $deg(gcd(x^n - 1, \sum_{i=0}^{n-1} \alpha^{q^i} x^{n-1-i})) = k$.

By analogy with the case of normal polynomials, a monic irreducible polynomial $P(x) \in \mathbb{F}_q[x]$ of degree n is called a k-normal polynomial (or N_k-polynomial) over $\mathbb{F}_q$ if its roots are k-normal elements of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$. Here, $P(x)$ has n distinct conjugate roots, of which $(n-k)$ are linearly independent. Recall that an element $\alpha \in \mathbb{F}_{q^n}$ is called a proper element of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$ if $\alpha \notin \mathbb{F}_{q^v}$ for any proper divisor v of n. So, the element $\alpha \in \mathbb{F}_{q^n}$ is a proper k-normal element of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$ if α is a k-normal and proper element of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$.

Using the above mention, a normal polynomial (or element) is a 0-normal polynomial (or element). Since the proper k-normal elements of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$ are the roots of a k-normal polynomial of degree n over $\mathbb{F}_q$, hence the k-normal polynomials of degree n over $\mathbb{F}_q$ is just another way of describing the proper k-normal elements of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$. Some results regarding constructions of spacial sequences of k-normal polynomials over $\mathbb{F}_q$, in the cases $k = 0$ and 1 can be found in [2, 4, 5, 10, 11] and [6], respectively.

In this paper, a recursive method for constructing the k-normal polynomials of higher degree from a given k-normal polynomial of degree n over $\mathbb{F}_q$ for each $0 \leq k \leq n-1$ is given.

2 Preliminary notes

We use the definitions, notations and results given by Huczynska [8], Gao [7] and Kyuregyan [10, 11], where similar problems are considered. We need the following results for our further study.

The trace of α in $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$, is given by $Tr_{\mathbb{F}_{q^n}|\mathbb{F}_q}(\alpha) = \sum_{i=0}^{n-1} \alpha^{q^i}$. For convenience, $Tr_{\mathbb{F}_{q^n}|\mathbb{F}_q}$ is denoted by $Tr_{q^n|q}$.

Proposition 2.1. *[6] Let $n = n_1 p^e$, for some $e \geq 1$ and $a, b \in \mathbb{F}_q^*$. Then the element α is a proper k-normal element of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$ if and only if $a + b\alpha$ is a proper k-normal element of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$.*

Let p denote the characteristic of $\mathbb{F}_q$ and let $n = n_1 p^e = n_1 t$, with $gcd(p, n_1) = 1$ and suppose that $x^n - 1$ has the following factorization in $\mathbb{F}_q[x]$:

$$x^n - 1 = (\varphi_1(x)\varphi_2(x) \cdots \varphi_r(x))^t, \tag{1}$$

where $\varphi_i(x) \in \mathbb{F}_q[x]$ are the distinct irreducible factors of $x^n - 1$. For each s, $0 \leq s < n$, let there is a $u_s > 0$ such that $R_{s,1}(x)$, $R_{s,2}(x)$, $\cdots$, $R_{s,u_s}(x)$ are all of the s degree polynomials dividing $x^n - 1$. So, from (1) we can write $R_{s,i}(x) = \prod_{j=1}^{r} \varphi_j{}^{t_{ij}}(x)$, for each $1 \leq i \leq u_s$, $0 \leq t_{ij} \leq t$. Let

$$\phi_{s,i}(x) = \frac{x^n - 1}{R_{s,i}(x)}, \tag{2}$$

for $1 \leq i \leq u_s$. Then, there is a useful characterization of the k-normal polynomials of degree n over $\mathbb{F}_q$ as follows.

Proposition 2.2. *[6] Let $F(x)$ be an irreducible polynomial of degree n over $\mathbb{F}_q$ and α be a root of it. Let $x^n - 1$ factor as (1) and let $\phi_{s,i}(x)$ be as in (2). Then $F(x)$ is a N_k-polynomial over $\mathbb{F}_q$ if and only if, there is j, $1 \leq j \leq u_k$, such that*

$$L_{\phi_{k,j}}(\alpha) = 0,$$

and also

$$L_{\phi_{s,i}}(\alpha) \neq 0,$$

for each s, $k < s < n$, and $1 \leq i \leq u_s$, where u_s is the number of all s degree polynomials dividing $x^n - 1$ and $L_{\phi_{s,i}}(x)$ is the linearized polynomial defined by

$$L_{\phi_{s,i}}(x) = \sum_{v=0}^{n-s} t_{iv} x^{q^v} \ \ if \ \ \phi_{s,i}(x) = \sum_{v=0}^{n-s} t_{iv} x^v.$$

The following propositions are useful for constructing N_k-polynomials over $\mathbb{F}_q$.

Proposition 2.3. *[3] Let $x^p - \delta_2 x + \delta_0$ and $x^p - \delta_2 x + \delta_1$ be relatively prime polynomials in $\mathbb{F}_q[x]$ and $P(x) = \sum_{i=0}^{n} c_i x^i$ be an irreducible polynomial of degree $n \geq 2$ over $\mathbb{F}_q$, and let $\delta_0, \delta_1 \in \mathbb{F}_q$, $\delta_2 \in \mathbb{F}_q^*$, $(\delta_0, \delta_1) \neq (0, 0)$. Then*

$$F(x) = (x^p - \delta_2 x + \delta_1)^n P\left(\frac{x^p - \delta_2 x + \delta_0}{x^p - \delta_2 x + \delta_1}\right)$$

is an irreducible polynomial of degree np over $\mathbb{F}_q$ if and only if $\delta_2{}^{\frac{q-1}{p-1}} = 1$ and

$$Tr_{q|p}\left(\frac{1}{A^p}\left((\delta_1 - \delta_0)\frac{P'(1)}{P(1)} - n\delta_1\right)\right) \neq 0,$$

where $A^{p-1} = \delta_2$, for some $A \in \mathbb{F}_q^*$.

Proposition 2.4. *[1] Let $x^p - x + \delta_0$ and $x^p - x + \delta_1$ be relatively prime polynomials in $\mathbb{F}_q[x]$ and let $P(x)$ be an irreducible polynomial of degree $n \geq 2$ over $\mathbb{F}_q$, and $0 \neq \delta_1, \delta_0 \in \mathbb{F}_p$, such that $\delta_0 \neq \delta_1$. Define*

$$F_0(x) = P(x)$$

$$F_k(x) = (x^p - x + \delta_1)^{t_{k-1}} F_{k-1}\left(\frac{x^p - x + \delta_0}{x^p - x + \delta_1}\right), \qquad k \geq 1$$

where $t_k = np^k$ denotes the degree of $F_k(x)$. Suppose that

$$Tr_{q|p}\left(\frac{(\delta_1 - \delta_0)F_0'(1) - n\delta_1 F_0(1)}{F_0(1)}\right) \cdot Tr_{q|p}\left(\frac{(\delta_1 - \delta_0)F_0'(\frac{\delta_0}{\delta_1}) + n\delta_1 F_0(\frac{\delta_0}{\delta_1})}{F_0(\frac{\delta_0}{\delta_1})}\right) \neq 0.$$

Then $(F_k(x))_{k\geq 0}$ is a sequence of irreducible polynomials over $\mathbb{F}_q$ of degree $t_k = np^k$, for every $k \geq 0$.

Lemma 2.5. *Let γ be a proper element of $\mathbb{F}_{q^n}$ and $\theta \in \mathbb{F}_p^*$, where $q = p^m$, $(m \in \mathbb{N})$. Then we have*

$$\sum_{j=0}^{p-1} \frac{1}{\gamma + j\theta} = -\frac{1}{\gamma^p - \gamma}. \tag{3}$$

Proof. By observing that

$$\sum_{j=0}^{p-1} \frac{1}{\gamma + j\theta} = \frac{1}{\gamma^p - \gamma}\left(\sum_{j=0}^{p-1} \frac{\gamma^p - \gamma}{\gamma + j\theta}\right), \tag{4}$$

it is enough to show that

$$\sum_{j=0}^{p-1} \frac{\gamma^p - \gamma}{\gamma + j\theta} = -1. \tag{5}$$

We note that

$$\begin{aligned}\sum_{j=0}^{p-1} \frac{\gamma^p - \gamma}{\gamma + j\theta} &= \sum_{j=0}^{p-1} \Big((\gamma + j\theta)^{p-1} - 1\Big) \\ &= \sum_{j=0}^{p-1} (\gamma + j\theta)^{p-1} \\ &= \sum_{j=1}^{p-1} \theta^j \binom{p-1}{j}\left(\sum_{i=1}^{p-1} i^j\right),\end{aligned} \tag{6}$$

where

$$\binom{p-1}{j} = \frac{(p-1)!}{(p-1-j)!j!}, \quad j \in \mathbb{F}_p^*.$$

On the other side, we know that

$$\sum_{i=1}^{p-1} i^j = \begin{cases} 0 \pmod p, & \text{if p-1} \nmid j \\ -1 \pmod p, & \text{if p-1} \mid j \end{cases} \tag{7}$$

and also $\theta^{p-1} = 1$. Thus by (6) and (7), the proof is completed.

□

3 Recursive construction N_k-polynomials

In this section we establish theorems which will show how propositions 2.2, 2.3 and 2.4 can be applied to produce infinite sequences of N_k-polynomials over $\mathbb{F}_q$. Recall that, the polynomial $P^*(x) = x^n P\left(\frac{1}{x}\right)$ is called the reciprocal polynomial of $P(x)$, where n is the degree of $P(x)$. In the case $k = 0$, some similar results of the following theorems has been obtained in [2], [4] and ([5], Theorems 3.3.1 and 3.4.1). We use of an analogous technique to that used in the above results, where similar problems are considered.

Theorem 3.1. *Let $P(x) = \sum_{i=0}^{n} c_i x^i$ be an N_k-polynomial of degree n over $\mathbb{F}_q$, for each $n = rp^e$, where $e \in \mathbb{N}$ and r equals 1 or is a prime different from p and q a primitive element modulo r. Suppose that $\delta \in \mathbb{F}_q^*$ and*

$$F(x) = (x^p - x + \delta)^n P^*\left(\frac{x^p - x}{x^p - x + \delta}\right). \tag{8}$$

Then $F^*(x)$ is an N_k-polynomial of degree np over $\mathbb{F}_q$ if $k < p^e$ and

$$Tr_{q|p}\left(\delta \frac{P^{*'}(1)}{P^*(1)}\right) \neq 0.$$

Proof. Since $P^*(x)$ is an irreducible polynomial over $\mathbb{F}_q$, so Proposition 2.3 and theorem's hypothesis imply that $F(x)$ is irreducible over $\mathbb{F}_q$.
Let $\alpha \in \mathbb{F}_{q^n}$ be a root of $P(x)$. Since $P(x)$ is an N_k-polynomial of degree n over $\mathbb{F}_q$ by theorem's hypothesis, then $\alpha \in \mathbb{F}_{q^n}$ is a proper k-normal element over $\mathbb{F}_q$.

Since q is a primitive modulo r, so in the case $r > 1$ the polynomial $x^{r-1} + \cdots + x + 1$ is irreducible over $\mathbb{F}_q$. Thus $x^n - 1$ has the following factorization in $\mathbb{F}_q[x]$:

$$x^n - 1 = (\varphi_1(x) \cdot \varphi_2(x))^t, \tag{9}$$

where $\varphi_1(x) = x - 1$, $\varphi_2(x) = x^{r-1} + \cdots + x + 1$ and $t = p^e$.

Letting that for each $0 \leq s < n$ and $1 \leq i \leq u_s$, $R_{s,i}(x)$ is the s degree polynomial dividing $x^n - 1$, where u_s is the number of all s degree polynomials dividing $x^n - 1$. So, from (9), we can write $R_{s,i}(x) = (x-1)^{s_{1,i}} \cdot (x^{r-1} + \cdots + x + 1)^{s_{2,i}}$, where $s = s_{1,i} + s_{2,i} \cdot (r-1)$ for each $0 \leq s_{1,i}, s_{2,i} \leq t$, except when $s_{1,i} = s_{2,i} = t$. So, we have

$$\phi_{s,i}(x) = \frac{x^n - 1}{R_{s,i}(x)} = \frac{x^n - 1}{(x-1)^{s_{1,i}} \cdot (x^{r-1} + \cdots + x + 1)^{s_{2,i}}} = \sum_{v=0}^{n-s} t_{s,i,v} x^v. \tag{10}$$

Since $P(x)$ is an N_k-polynomial of degree n over $\mathbb{F}_q$, so by Proposition 2.2, there is a j, $1 \leq j \leq u_k$, such that

$$L_{\phi_{k,j}}(\alpha) = 0, \tag{11}$$

and also

$$L_{\phi_{s,i}}(\alpha) \neq 0, \tag{12}$$

for each $k < s < n$ and $1 \leq i \leq u_s$. Further, we proceed by proving that $F^*(x)$ is a k-normal polynomial. Let α_1 be a root of $F(x)$. Then $\beta_1 = \frac{1}{\alpha_1}$ is a root of its reciprocal polynomial $F^*(x)$. Note that by (9), the polynomial $x^{np} - 1$ has the following factorization in $\mathbb{F}_q[x]$:

$$x^{np} - 1 = (\varphi_1(x) \cdot \varphi_2(x))^{pt}, \tag{13}$$

where $\varphi_1(x) = x - 1$, $\varphi_2(x) = x^{r-1} + \cdots + x + 1$ and $t = p^e$.

Letting that for each $0 \leq s' < np$ and $1 \leq i' \leq u'_{s'}$, $R'_{s',i'}(x)$ is the s' degree polynomial dividing $x^{np} - 1$, where $u'_{s'}$ is the number of all s' degree polynomials dividing $x^{np} - 1$. So, from

(13) we can write $R'_{s',i'}(x) = (x-1)^{s'_{1,i'}} \cdot (x^{r-1} + \cdots + x + 1)^{s'_{2,i'}}$, where $s' = s'_{1,i'} + s'_{2,i'} \cdot (r-1)$ for each $0 \leq s'_{1,i'}, s'_{2,i'} \leq pt$, except when $s'_{1,i'} = s'_{2,i'} = pt$. Therefore by considering

$$H'_{s',i'}(x) = \frac{x^{np}-1}{R'_{s',i'}(x)}, \tag{14}$$

and Proposition 2.2, $F^*(x)$ is an N_k-polynomial of degree np over $\mathbb{F}_q$ if and only if there is a j', $1 \leq j' \leq u'_k$, such that

$$L_{H'_{k,j'}}(\beta_1) = 0,$$

and also

$$L_{H'_{s',i'}}(\beta_1) \neq 0,$$

for each $k < s' < np$ and $1 \leq i' \leq u'_{s'}$. Consider

$$\begin{aligned} H_{s,i}(x) &= \frac{x^{np}-1}{R_{s,i}(x)} \\ &= \frac{x^n-1}{R_{s,i}(x)} \left(\sum_{j=0}^{p-1} x^{jn} \right), \end{aligned} \tag{15}$$

for each $0 \leq s < n$ and $1 \leq i \leq u_s$. By (10) we obtain

$$H_{s,i}(x) = \phi_{s,i}(x) \left(\sum_{j=0}^{p-1} x^{jn} \right) = \sum_{v=0}^{n-s} t_{s,i,v} \left(\sum_{j=0}^{p-1} x^{jn+v} \right).$$

It follows that

$$L_{H_{s,i}}(\beta_1) = \sum_{v=0}^{n-s} t_{s,i,v} \left(\sum_{j=0}^{p-1} (\beta_1)^{p^{jmn}} \right)^{p^{mv}},$$

or

$$L_{H_{s,i}}(\beta_1) = L_{H_{s,i}}\left(\frac{1}{\alpha_1}\right) = \sum_{v=0}^{n-s} t_{s,i,v} \left(\sum_{j=0}^{p-1} \left(\frac{1}{\alpha_1}\right)^{p^{jmn}} \right)^{p^{mv}}. \tag{16}$$

From (8), if α_1 is a zero of $F(x)$, then $\frac{{\alpha_1}^p - \alpha_1 + \delta}{{\alpha_1}^p - \alpha_1}$ is a zero of $P(x)$, and therefore it may assume that

$$\alpha = \frac{{\alpha_1}^p - \alpha_1 + \delta}{{\alpha_1}^p - \alpha_1},$$

or

$$\frac{\alpha - 1}{\delta} = ({\alpha_1}^p - \alpha_1)^{-1}. \tag{17}$$

Now, by (17) and observing that $P(x)$ is an irreducible polynomial of degree n over $\mathbb{F}_q$, we obtain

$$\frac{\alpha - 1}{\delta} = \left(\frac{\alpha - 1}{\delta}\right)^{p^{mn}} = ({\alpha_1}^{p^{mn+1}} - {\alpha_1}^{p^{mn}})^{-1}. \tag{18}$$

It follows from (17) and (18) that

$$({\alpha_1}^{p^{mn+1}} - {\alpha_1}^{p^{mn}})^{-1} = ({\alpha_1}^p - \alpha_1)^{-1}. \tag{19}$$

Also observing that $F(x)$ is an irreducible polynomial of degree np over $\mathbb{F}_q$, we have $({\alpha_1}^p - \alpha_1) \neq 0$ and $({\alpha_1}^{p^{mn+1}} - {\alpha_1}^{p^{mn}}) \neq 0$. Hence by (19)

$$({\alpha_1}^{p^{mn}} - \alpha_1)^p = ({\alpha_1}^{p^{mn}} - \alpha_1). \tag{20}$$

It follows from (20) that ${\alpha_1}^{p^{mn}} - \alpha_1 = \theta \in \mathbb{F}_p^*$. Hence ${\alpha_1}^{p^{mn}} = \alpha_1 + \theta$ and

$${\alpha_1}^{p^{2mn}} = (\alpha_1 + \theta)^{p^{mn}} = {\alpha_1}^{p^{mn}} + \theta^{p^{mn}} = \alpha_1 + \theta + \theta = \alpha_1 + 2\theta.$$

It is easy to show that

$${\alpha_1}^{p^{jmn}} = \alpha_1 + j\theta, \quad for \;\; 1 \le j \le p-1,$$

or

$$\left(\frac{1}{\alpha_1}\right)^{p^{jmn}} = \frac{1}{\alpha_1 + j\theta}, \quad for \;\; 1 \le j \le p-1. \tag{21}$$

From (16) and (21), we immediately obtain

$$L_{H_{s,i}}(\beta_1) = \sum_{v=0}^{n-s} t_{s,i,v} \left(\sum_{j=0}^{p-1} \frac{1}{\alpha_1 + j\theta} \right)^{p^{mv}}. \tag{22}$$

Thus, by (17), (22) and Lemma 2.5 we have

$$\begin{aligned} L_{H_{s,i}}(\beta_1) &= \sum_{v=0}^{n-s} t_{s,i,v} \left(-\frac{1}{\alpha_1^p - \alpha_1} \right)^{p^{mv}} \\ &= \frac{1}{\delta} \sum_{v=0}^{n-s} t_{s,i,v} (1-\alpha)^{p^{mv}} \\ &= L_{\phi_{s,i}} \left(\frac{1-\alpha}{\delta} \right). \end{aligned} \tag{23}$$

Since α is a zero of $P(x)$, then α will be a k-normal element in $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$. Thus according to Proposition 2.1, the element $\frac{1-\alpha}{\delta}$ will also be a k-normal element.

since $\frac{1-\alpha}{\delta}$ is a root of $P(-\delta x + 1)$, so by (23) and Proposition 2.2, there is a j, $1 \le j \le u_k$, such that

$$L_{H_{k,j}}(\beta_1) = 0,$$

and also

$$L_{H_{s,i}}(\beta_1) \neq 0,$$

for each s, $k < s < n$ and $1 \le i \le u_s$. So, there is a j', $1 \le j' \le u'_k$, such that, $L_{H'_{k,j'}}(\beta_1) = L_{H_{k,j}}(\beta_1) = 0$. On the other side, by (14) and (15), for each s', $k < s' < np$ and $1 \le i' \le u'_{s'}$, there is s, $k < s < n$ and $1 \le i \le u_s$ such that $H'_{s',i'}(x)$ divide $H_{s,i}(x)$. It follows that

$$L_{H'_{s',i'}}(\beta_1) \neq 0,$$

for each s', $k < s' < np$ and $1 \le i' \le u'_{s'}$. The proof is completed. □

In the following theorem, a computationally simple and explicit recurrent method for constructing higher degree N_k-polynomials over $\mathbb{F}_q$ starting from an N_k-polynomial is described.

Theorem 3.2. *Let $P(x)$ be an N_k-polynomial of degree n over $\mathbb{F}_q$, for each $n = rp^e$, where $e \in \mathbb{N}$ and r equals 1 or is a prime different from p and q a primitive element modulo r. Define*

$$\begin{aligned} F_0(x) &= P^*(x) \\ F_u(x) &= (x^p - x + \delta)^{np^{u-1}} F_{u-1}\left(\frac{x^p - x}{x^p - x + \delta} \right), \end{aligned} \tag{24}$$

where $\delta \in \mathbb{F}_p^*$. *Then* $(F_u^*(x))_{u\geq 0}$ *is a sequence of* N_k*-polynomials of degree* np^u *over* $\mathbb{F}_q$ *if* $k < p^e$ *and*

$$Tr_{q|p}\left(\frac{P^{*'}(0)}{P^*(0)}\right) \cdot Tr_{q|p}\left(\frac{P^{*'}(1)}{P^*(1)}\right) \neq 0,$$

where $P^{*'}(0)$ *and* $P^{*'}(1)$ *are the formal derivative of* $P^*(x)$ *at the points* $x = 0$ *and* $x = 1$, *respectively.*

Proof. By Proposition 2.4 and hypotheses of theorem for each $u \geq 1$, $F_u(x)$ is an irreducible polynomial over $\mathbb{F}_q$. Consequently, $(F_u^*(x))_{u\geq 0}$ is a sequence of irreducible polynomials over $\mathbb{F}_q$. The proof of k-normality of the irreducible polynomials $F_u^*(x)$, for each $u \geq 1$ is implemented by mathematical induction on u. In the case $u = 1$, by Theorem 3.1 ${F_1}^*(x)$ is a k-normal polynomial.

For $u = 2$ we show that ${F_2}^*(x)$ is also a k-normal polynomial. To this end we need to show that the hypothesis of Theorem 3.1 are satisfied. By Theorem 3.1, ${F_2}^*(x)$ is a k-normal polynomial if

$$Tr_{q|p}\left(\frac{F_1'(1)}{F_1(1)}\right) \neq 0,$$

since $\delta \in \mathbb{F}_p^*$. We apply (24) to compute

$$F_u(0) = F_u(1) = \delta^{un} P^*(0), \quad u = 1, 2, \ldots. \tag{25}$$

We calculate the formal derivative of $F_1'(x)$ at the points $x = 0$ and $x = 1$. According to (24) the first derivative of $F_1(x)$ is

$$\begin{aligned}
{F_1}'(x) &= -n(x^p - x + \delta)^{n-1} {F_0}'\left(\frac{x^p - x}{x^p - x + \delta}\right) \\
&\quad + (x^p - x + \delta)^n \cdot \left(\frac{(px^{p-1} - 1)(x^p - x + \delta) - (px^{p-1} - 1)(x^p - x)}{(x^p - x + \delta)^2}\right) \\
&\quad \cdot {F_0}'\left(\frac{x^p - x}{x^p - x + \delta}\right) \\
&= -\delta(x^p - x + \delta)^{n-2} \cdot {P^*}'\left(\frac{x^p - x}{x^p - x + \delta}\right),
\end{aligned}$$

and at the points $x = 0$ and $x = 1$

$${F_1}'(0) = -{F_1}'(1) = -\delta^{n-1} {P^*}'(0) \tag{26}$$

which is not equal to zero by the condition $Tr_{q|p}\left(\frac{P^{*'}(0)}{P^*(0)}\right) \neq 0$ in the hypothesis of theorem, since $\delta \in \mathbb{F}_p^*$. From (26) and (25)

$$\begin{aligned}
Tr_{q|p}\left(\frac{F_1'(1)}{F_1(1)}\right) &= Tr_{q|p}\left(\frac{-\delta^{n-1} P^{*'}(0)}{\delta^n P^*(0)}\right) \\
&= -\frac{1}{\delta} Tr_{q|p}\left(\frac{P^{*'}(0)}{P^*(0)}\right),
\end{aligned} \tag{27}$$

which is not equal to zero by hypothesis of theorem. Hence the polynomial $F_2^*(x)$ is a k-normal polynomial. If induction holds for $u - 1$, then it must hold also for u, that is by assuming that $F_{u-1}^*(x)$ is a k-normal polynomial, we show that $F_u^*(x)$ is also a k-normal polynomial.
Let $u \geq 3$. By Theorem 3.1, $F_u^*(x)$ is a k-normal polynomial if

$$Tr_{q|p}\left(\frac{F_{u-1}'(1)}{F_{u-1}(1)}\right) \neq 0,$$

since $\delta \in \mathbb{F}_p^*$. We calculate the formal derivative of $F'_{u-1}(x)$ at the points 1 and 0. By (24) the first derivative of $F_{u-1}(x)$ is

$$\begin{aligned} F'_{u-1}(x) &= (x^p - x + \delta)^{np^{u-2}} \left(\frac{(px^{p-1} - 1)(x^p - x + \delta) - (px^{p-1} - 1)(x^p - x)}{(x^p - x + \delta)^2} \right) \\ &\quad \cdot F'_{u-2} \left(\frac{x^p - x}{x^p - x + \delta} \right) \\ &= -\delta (x^p - x + \delta)^{np^{u-2}-2} F'_{u-2} \left(\frac{x^p - x}{x^p - x + \delta} \right), \end{aligned}$$

and at the point $x = 0$ and $x = 1$

$$F'_{u-1}(0) = F'_{u-1}(1) = -\delta^{np^{u-2}-1} F'_{u-2}(0) = -\delta^{n-1} F'_{u-2}(0). \tag{28}$$

So we have

$$F'_{u-1}(0) = F'_{u-1}(1) = (-1)^{u-2} \delta^{(n-1)(u-2)} F'_1(0), \tag{29}$$

which is not equal to zero by (26) and the condition $Tr_{q|p}\left(\frac{P^{*'}(0)}{P^*(0)}\right) \neq 0$ in the hypothesis of theorem, since $\delta \in \mathbb{F}_p^*$. Also

$$Tr_{q|p}\left(\frac{F'_{u-1}(1)}{F_{u-1}(1)}\right) = (-1)^{u-2} \frac{1}{\delta^{(u-2)}} Tr_{q|p}\left(\frac{F'_1(1)}{F_1(1)}\right),$$

which is not equal to zero by (27) and hypothesis of theorem. The theorem is proved. □

References

[1] S. Abrahamyan, M. K. Kyureghyan, A recurrent method for constructing irreducible polynomials over finite fields, proceeding of the 13th international conference on computer algebra in scientific computing, 2011, 1-9.

[2] S. Abrahamyan, M. K. Kyureghyan, New recursive construction of normal polynomials over finite fields, proceeding of the 11th international conference on finite fields and their applications, 2013, 1-10.

[3] S. Abrahamyan, M. Alizadeh, M. K. Kyureghyan, Recursive constructions of irreducible polynomials over finite fields, Finite Fields Appl., 18 (2012), 738-745.

[4] M. Alizadeh, S. Abrahamyan, S. Mehrabi, M. K. Kyuregyan, Constructing of N-polynomials over finite fields, International Journal of Algebra, Vol. 5, No. 29, 2011, 1437-1442.

[5] M. Alizadeh, Construction of Irreducible and Normal Polynomials over Finite Fields, Ph.D. Thesis, National Academy of Sciences of Armenia, Yerevan, 2013.

[6] M. Alizadeh, Some notes on the k-normal elements and k-normal polynomials over finite fields, J. Algebra Appl., DOI:10.1142/S0219498817500062, 2016.

[7] S. Gao, Normal bases over finite fields, Ph.D. Thesis, Waterloo, 1993.

[8] S. Huczynska, G. L. Mullen, D. Panario and D. Thomson, Existence and properties of k-normal elements over finite fields, Finite Fields Appl., 24 (2013), 170-183.

[9] D. Jungnickel, Finite Fields: Structure and Arithmetics, Wissenschaftsverlag, (Mannheim) 1993.

[10] M. K. Kyuregyan, Iterated construction of irreducible polynomials over finite fields with linearly independent roots, Finite Fields Appl., 10 (2004) 323-341.

[11] M. K. Kyuregyan, Recursive construction of N-polynomials over $GF(2^s)$, Discrete Applied Mathematics, No. 156, 2008, 1554-1559.

[12] H. W. Lenstra and R. Schoof, Primitive normal bases for finite fields, Mathematics of Computation, 48 (1987), 217-231.

[13] S. Schwartz, Irreducible polynomials over finite fields with linearly independent roots, Math. Slovaca, Vol. 38, 1988, 147-158.